

From Bench to Burden: Illuminating the Unseen Challenges in Pakistani Healthcare

Inayat Ullah

Department of Internal Medicine, Lady Reading Hospital, Peshawar, KPK, Pakistan

***Corresponding Author:**

Email: inayatkmc@yahoo.com

Article History

Received: 17-11-2025

Revised: 21-12-2025

Accepted: 28-12-2025

Published: 30-12-2025

How to cite

Inayat Ullah. From bench to burden: illuminating the unseen challenges in Pakistani healthcare. Pak J Clin Res. 2025;2(2):17-18. doi:10.65761/pjcr.2025.2.2.30.

DOI: <https://doi.org/10.65761/pjcr.2025.2.2.30>

Keywords: Medical Students; Musculoskeletal Pain; Hypoglycemia; Neonatal Care; Cerebellopontine Angle; Surgical Outcomes; Pakistan

Clinical research often gravitates toward dramatic therapeutic breakthroughs and high-technology interventions. While those pursuits are essential, there exists a quieter, equally critical domain: the systematic study of everyday suffering, vulnerable populations, and the hidden burdens carried by both patients and caregivers. The three original articles comprising Volume 2, Issue 2 of the Pakistan Journal of Clinical Research (PJCR) do precisely this. They turn our gaze away from the operating theater's brilliance and toward the aching backs of future doctors, the fragile glucose homeostasis of our smallest citizens, and the comparative nuances of two common but distinct intracranial tumors. Collectively, they remind us that great clinical science is measured not only in survival curves but in the relief of pervasive, often overlooked distress.

The Silent Epidemic Among Tomorrow's Healers

We begin with a study that should give every medical educator pause. Arshad and colleagues, in their cross-sectional analysis of medical students in Peshawar, document a significant association between musculoskeletal pain and gastrointestinal symptoms among this already stretched population. The finding is more than a statistical curiosity. It is a window into the physical toll of a demanding training culture—long hours of static postures, irregular meals, chronic stress, and neglect of personal health. The gut-brain axis is no longer a theoretical concept; here it manifests as real, measurable comorbidity. This research compels us to ask an uncomfortable question: If we are training students to

care for patients' bodies, who is caring for theirs? The authors have provided evidence that fatigue, lumbar pain, and dyspepsia are not incidental but intertwined. Medical curricula across Pakistan must now consider structured wellness interventions not as luxuries but as prerequisites for clinical competence.

Protecting the Most Fragile: Hypoglycemia in Preterm Neonates

From young adults we turn to the most vulnerable among us—preterm infants. Shagufta Shehzadi and Mehwish Nazir present a sobering prevalence estimate of hypoglycemia in a neonatal intensive care unit (NICU). Preterm infants, by virtue of immature metabolic pathways and limited glycogen stores, exist on a knife's edge. Undetected or inadequately managed hypoglycemia remains a preventable cause of neurodevelopmental harm. This study is a necessary audit of current practice. It does not merely report numbers; it issues a quality-of-care indicator. The prevalence identified calls for standardized screening protocols, tighter glucose monitoring, and perhaps most importantly, a culture of zero-tolerance for metabolic instability in the NICU. Every percentage point matter when the outcome is a developing brain. The authors deserve credit for placing this issue squarely on our collective table.

Surgical Nuance in the Cerebellopontine Angle

The third article, by Asif Khan and colleagues, offers a retrospective comparison of surgical outcomes for schwannomas versus meningiomas in the



cerebellopontine angle-a notoriously challenging anatomical region. While both tumors are benign, their behavior, adherence to neurovascular structures, and surgical morbidity differ substantially. This study provides Pakistani-specific data to guide preoperative counseling and surgical planning. The retrosigmoid approach, as examined here, yields valuable insights into expected functional recovery, complication rates, and resection completeness. For neurosurgeons practicing in resource-variable settings, such comparative effectiveness research is gold. It allows for informed decisions about case selection, surgical aggressiveness, and postoperative care pathways. We encourage the authors and their peers to extend this work into prospective registries, which would further strengthen evidence for surgical decision-making in Pakistan.

A Unified Editorial Vision

Three articles, three distinct domains-yet a common thread binds them: the primacy of patient-centered and caregiver-centered outcomes. Whether we study a medical student's chronic pain, a preterm infant's glucose level, or a postoperative cranial nerve function, the underlying question is the same: Are we truly seeing the full burden of illness and risk?

PJCR remains committed to publishing research that answers this question with honesty and precision. Our peer-review process prioritizes methodological soundness, ethical integrity, and clinical relevance. We continue to uphold the standards of COPE and ICMJE, and we are pleased that all published work in this issue carries Crossref DOIs and is discoverable via Scilit. These technical foundations are not ending in themselves but means to a greater end: ensuring that high-quality evidence from Pakistan reaches clinicians, policymakers, and educators both nationally and globally.

Looking Ahead

We invite submissions that challenge clinical dogma, illuminate neglected populations, and bridge disciplinary divides. Particularly welcome are studies on medical trainee health, neonatal and maternal outcomes, comparative surgical effectiveness, and cross-specialty syndromes that defy simple classification. We also reaffirm our support for early-career investigators-many of whom are the first authors in this issue-whose fresh perspectives and local insights are the lifeblood of contextualized research.

In closing, we thank our authors, reviewers, readers, and the editorial team. This issue, though small in number of articles, is large in its implications. It tells us that healing begins not with grand gestures but with the careful, compassionate measurement of pain, risk, and resilience.

Ethical Approval

Ethical approval was not required for this study as it did not involve human or animal participants.

Data Availability

Data sharing is not applicable to this article as it is an editorial piece.

Author Contributions

Inayat Ullah contributed to the conceptualization, methodology, formal analysis, original draft writing, and review and editing of the manuscript.

Informed Consent

Informed consent was not required for this study as it does not involve human participants or patient-specific data.

Funding

This research received no external funding.

Conflict of Interest

The authors declare that they have no competing interests.

Acknowledgments

The author acknowledges the editorial team of the Pakistan Journal of Clinical Research (PJCR) and the colleagues at Lady Reading Hospital, Peshawar, for their continued academic support and encouragement.

References

1. Huma S, Mahar SA, Kalwar S, Ahmed J, Saeed U, Shah SSH. Impact of chronic musculoskeletal pain on gastrointestinal symptoms and mental health outcomes in adult Pakistanis. *MedERA*. 2025;7(1). doi:10.61982/medera.v7i1.248.
2. Bashir NU, Bibi S, Adnan MS, Gilani SYH. Frequency and pattern of hypoglycaemia in low birth weight babies during first 48 hours of life. *Pak J Physiol*. 2025;21(3):11-13. doi:10.69656/pjp.v21i3.1852.
3. Khokhar TI, Ghafoor A, Shakir M, Khan FI, Qadri HM. Comparative analysis of clinical outcomes of cerebellopontine angle meningioma versus cerebellopontine angle schwannoma: a retrospective insight from Pakistan. *Pak J Med Sci*. 2025;41(13(PINS-NNOS)):S43-S48. doi:10.12669/pjms.41.13(PINS-NNOS).13374.
4. Sadia H, et al. Enhancing medication appropriateness: insights from the STOPP criteria version 3 on prescribing practices among older adults in Pakistan. *Front Pharmacol*. 2025;16:1551819. doi:10.3389/fphar.2025.1551819.